

Barnsley Self-Care Guidance

Guidance on conditions for which over the counter items should not routinely be prescribed in primary care

The full NHS England guidance is available at: <https://www.england.nhs.uk/wp-content/uploads/2018/03/otc-guidance-for-ccgs.pdf>

Barnsley CCG has made the decision to adopt all of the recommendations within the NHSE guidance. This document has been produced to support primary care clinicians with implementation of the recommendations in practice. It can also be used by community pharmacists to help provide information to the patient on self-care for each condition and to help signpost the patient to the right service.

The guidance is intended to encourage people to self-care for minor illnesses as the first stage of treatment. It is envisioned that in most cases (unless specified) these minor conditions will clear up with appropriate self-care. If symptoms are not improving or responding to treatment, then patients should be encouraged to seek further advice.

The guidance is not intended to discourage patients from going to the GP when it is appropriate to do so.

This guidance applies to all patients, including those who would be exempt from paying prescription charges, unless they fall under the exceptions outlined below (general exceptions) or within the table below under each condition/item.

GPs and/or pharmacists should refer patients to NHS UK, the Self Care Forum or NHS 111 for further advice on when they should seek GP Care.

Why has NHSE developed this guidance?

In the year prior to June 2017, the NHS spent approximately £569 million on prescriptions for medicines, which could otherwise be purchased over the counter (OTC) from a pharmacy and/or other outlets such as petrol stations or supermarkets. In the 12 months prior to June 2017, NHS Barnsley spent approximately £1.4 million on prescriptions for medicines, which could otherwise be purchased OTC.

These prescriptions include items for a condition:

- That is considered to be self-limiting and so does not need treatment as it will heal or be cured of its own accord;
- Which lends itself to self-care i.e. the person suffering does not normally need to seek medical advice and can manage the condition by purchasing OTC items directly.

These prescriptions also include other common items:

- That can be purchased over the counter, sometimes at a lower cost than that which would be incurred by the NHS;
- For which there is little evidence of clinical effectiveness.

A wide range of information is available to the public on the subjects of health promotion and the management of minor self-treatable illnesses. Advice from organisations such as the [Self Care Forum](#) and [NHS.uk](#) website is readily available on the internet. Many community pharmacies are also open extended hours including weekends and are ideally placed to offer advice on the management of minor conditions and lifestyle interventions. [The Royal Pharmaceutical Society](#) offers advice on over the counter products that should be kept in a medicine cabinet at home to help patients treat a range of self-treatable illnesses.

General exceptions to the guidance

There are certain scenarios where patients should continue to have their treatments prescribed and these are outlined below.

To note that for vitamins, minerals, probiotics and those self-limiting conditions where there is limited evidence of clinical effectiveness for the treatments used (e.g. OTC items for cough, sore throat and infant colic), then the **general exceptions do not apply**. Specific exceptions are included in the table below under each item/condition.

- Patients prescribed an OTC treatment for a **long term condition** (e.g. regular pain relief for chronic arthritis or treatments for inflammatory bowel disease).
- For the treatment of more **complex forms of minor illnesses** (e.g. severe migraines that are unresponsive to over the counter medicines).
- For those patients that have symptoms that suggest the condition is not minor (i.e. those with **red flag symptoms**, for example indigestion with very bad pain).
- Treatment for **complex patients** (e.g. immunosuppressed patients).
- Patients on **prescription only treatments**.
- Patients prescribed OTC products to treat an **adverse effect or symptom of a more complex illness** and/or prescription only medications should continue to have these products prescribed on the NHS.
- Circumstances where the **product licence** doesn't allow the product to be sold over the counter to certain groups of patients. This may vary by medicine, but could include babies, children and/or women who are pregnant or breast-feeding. Community Pharmacists will be aware of what these are and can advise accordingly. More information on over the counter restrictions is available in the table below and in the [PrescQIPP bulletin 227: Over the counter items - GP guide to self care](#).
- Patients with a minor condition suitable for self-care that has **not responded sufficiently to treatment** with an OTC product.
- Patients where the clinician considers that the presenting symptom is due to a **condition that would not be considered a minor condition**.
- Circumstances where the prescriber believes that in their **clinical judgement**, exceptional circumstances exist that warrant deviation from the recommendation to self-care.
- Individual patients where the clinician considers that their ability to self-manage is compromised as a consequence of medical, mental health or significant social vulnerability to the extent that **their health and/or wellbeing could be adversely affected, if reliant on self-care**. To note that being exempt from paying a prescription charge does not automatically warrant an exception to the guidance. Consideration should also be given to safeguarding issues.

Implementing the NHS England guidance in care settings.

The [CQC](#) supports the principle of promoting self-care and the implementation of the NHS England OTC guidance in social care settings, provided appropriate safeguards are in place.

It is important that implementing the OTC guidance does not restrict safe access to necessary medicines for vulnerable people, such as the elderly and people with learning difficulties, living in care homes, assisted living environments, or receiving social care services. Before stopping prescribing for the conditions included in the guidance and recommending that a care home resident purchases their own medicine the clinician should consider the following on an individual patient basis:

1. NHSE General Exceptions (see [page 2](#))

2. The Care Providers' Medication Policies

The CQC recommends that care providers should have a policy in place to allow access to OTC medicines in appropriate situations. This is different to the use of homely remedies.

Clinicians should liaise closely with carers before recommending the use of OTC medication in any care setting to ensure that:

- The patient is able to obtain the medicine in a safe and timely manner
- The patient is able to self-administer safely, or,
- If carers will be required to administer the medicine, that this is permitted within the care providers own medication policy.

After agreement with the care provider a clinician may recommend a resident, relative or care staff purchases an OTC medicine. The clinician should give verbal or written instructions on the use of the medicine to the patient and carers. This must be recorded in the individual care plan.

3. Use of Homely Remedies

In some cases a care home resident may be able to make use of the care home's own homely medicines. Homely medicines are a range of medicines that a care home may choose to purchase and keep in stock for use by their residents to treat minor conditions. They are the equivalent of the OTC medicines that would commonly be kept in any household, for example paracetamol for a headache or an antacid for indigestion. Use of homely medicines has to be pre-approved by a suitable health care professional e.g. doctor or pharmacist, in advance of use or at the time of need; and a record kept in the residents care plan. Their approval should be reviewed regularly. Homely medicines can only be taken for a short period of time without consulting a doctor, usually a maximum of 48 hours. Although the CQC supports the use of homely medicines, their availability is dependent on the individual care provider's medication policy.

Further information on implementing the NHS England guidance in care settings can be found at: <https://www.england.nhs.uk/medicines/conditions-for-which-over-the-counter-items-should-not-routinely-be-prescribed/conditions-for-which-over-the-counter-items-should-not-routinely-be-prescribed-in-primary-care-implementation-resources/faqs/>

Suggested actions for prescribers

New patients

- If a patient presents with a condition that can be managed with an over the counter preparation and / or self-care, sign post the patient to a pharmacy who will be able to offer advice and recommend a suitable preparation (see the [useful resources](#) section for 'no prescription required' leaflets and the patient information resources included in the following tables).
- Promote the self-care websites ([NHS.uk](#), [Self-Care Forum](#)) and the clinical expertise of community pharmacies in supporting management of minor ailments and self-limiting conditions.

Use the NHS Barnsley CCG 'Prescribing of over the counter medicines is changing' leaflets and posters to explain the changes to prescribing of over the counter medicines and why it is now recommended that over the counter medicines are purchased. These can be obtained from the Medicines Management Team.

Existing patients

Review all patients prescribed a medicine for a condition in the tables below. Determine whether to:

- Stop prescribing the medicine and provide the patient with advice on self-care or non-drug alternatives. For items available to purchase over the counter, recommend self-care and purchase of the medicine with support and advice from the community pharmacist wherever appropriate (note some preparations will only be available through community pharmacies – 'P' medicines).
- If it is appropriate in 'exceptional circumstances' to continue treatment, ensure that the patient is assessed and reviewed as appropriate. Document the exception/reason for prescribing in the patients notes.

Consider using patient and professional resources to explain the changes to prescribing of over the counter medicines and why it is now recommended that over the counter medicines are purchased. NHS Barnsley CCG 'Prescribing of over the counter medicines is changing' leaflets and posters can be obtained from the Medicines Management Team. Also see the [useful resources](#) section for 'no prescription required' leaflets and the patient information resources included in the following tables.

The tables on the following pages list the conditions/items where prescribing should be restricted, exceptions when prescribing may be appropriate, examples of products available to purchase, over the counter restrictions and links to self-care leaflets and advice.

Links to information and resources for each condition have also been included within the tables to help support clinicians with the decision of whether prescribing may be appropriate for each condition/item in the guidance (e.g. information on 'red flag symptoms'). However this information is not exhaustive and the decision of whether to prescribe remains the clinical decision of the clinician.

Condition/item for which prescribing should be restricted	
Items of limited clinical effectiveness	Minor conditions suitable for self-care continued
Probiotics page 6	Mild to moderate hay fever/seasonal rhinitis page 14
Vitamins and minerals page 6	Minor burns and scalds page 15
Self-limiting conditions	Minor conditions associated with pain, discomfort and/fever. (e.g. aches and sprains, headache, period pain, back pain) page 16
Acute sore throat page 7	Mouth ulcers page 16
Cold sores page 8	Nappy rash page 17
Conjunctivitis page 8	Oral thrush page 17
Coughs and colds and nasal congestion page 8	Prevention of dental caries page 17
Cradle cap page 9	Ringworm/ athlete's foot page 17
Haemorrhoids page 9	Teething/mild toothache page 17
Infant colic page 9	Threadworms page 18
Mild cystitis page 9	Travel Sickness page 18
Minor conditions suitable for self-care	Warts and Verrucae page 18
Mild Irritant Dermatitis page 10	
Dandruff page 10	
Diarrhoea (adults) page 10	
Dry eyes/sore tired eyes page 10	
Earwax page 11	
Excessive sweating (hyperhidrosis) page 11	
Head lice page 11	
Indigestion and heartburn page 12	
Infrequent constipation page 12	
Infrequent migraine page 13	
Insect bites and stings page 13	
Mild acne page 13	
Mild dry skin page 14	
Sunburn page 14	
Sun protection page 14	

Condition/item for which prescribing should be restricted	Exceptions to the guidance when prescribing may be appropriate	Examples of products available to purchase (note: some of the products may be pharmacy only). This is not an exhaustive list.	Over the counter restrictions*	Information and resources for the patient (key self-care messages, signposting)	Information and resources for the clinician
Items of limited clinical effectiveness					
Probiotics	<p>No exceptions. There is currently insufficient clinical evidence to support prescribing of probiotics within the NHS for the treatment or prevention of diarrhoeas of any cause.</p> <p>All probiotics are classed as food supplements</p>	Probiotic sachets	N/A – all probiotics are classed as food supplements		<p>Public Health England C. Difficile guidance</p> <p>NICE CG 84: Diarrhoea and vomiting caused by gastroenteritis in under 5s: diagnosis and management</p>
Vitamins and minerals	<p>Medically diagnosed deficiency, including for those patients who may have a lifelong or chronic condition or have undergone surgery that results in malabsorption. Continuing need should however be reviewed on a regular basis.</p> <p><i>NB maintenance or preventative treatment is not an exception.</i></p> <p>Calcium and vitamin D for osteoporosis.</p> <p>Malnutrition including alcoholism (see NICE Guidance)</p>	<p>Vitamin D (NB. Not the high strength preparations used to treat deficiency), <i>The clinician should advise the patient which strength of vitamin D to purchase. Barnsley vitamin D guidelines are available on the BEST website:</i></p> <p>Management of Low Vitamin D Level: A Barnsley Guideline (Adults)</p> <p>Guidelines for the management of children with suspected vitamin D deficiency in primary care setting</p>	Many are food supplements so limited information available	<p>Do I need vitamin supplements? - NHS</p> <p>Healthy Start vitamins</p> <p>NHS Choices: Supplements, Who Needs Them? A behind the Headlines Report, June 2011</p>	<p>PrescQIPP bulletin 107, August 2015: the prescribing of vitamins and minerals including vitamin B preparations (DROP-list)</p> <p>PHE publishes new advice on vitamin D - GOV.UK</p>

	<p>Vitamin supplements for premature and low birth weight babies (up to one year of age) as advised by the hospital.</p> <p><i>Patients suitable to receive Healthy start vitamins for pregnancy or children between the ages 6 months to their fourth birthday. (NB this is not on prescription but commissioned separately. See Barnsley Healthy Start Vitamin D Guidance - Children)</i></p> <p>This guidance does not apply to prescription only vitamin D analogues such as alfacalcidol and these should continue to be prescribed.</p>	<p>Vitamin B compound tablets</p> <p>Vitamin C effervescent 1g tablets</p> <p>Multivitamin preparations</p>			
Self-limiting conditions					
Acute sore throat	'Red flag' symptoms	<p>Benzydamine for example as Difflam® Spray and Difflam® Oral Rinse</p> <p>Lozenges for example Strepsils®, Throaties® and Tyrozets®</p> <p>Throat sprays for example Ultra Chloraseptic®</p>	<p>All preparations – pregnancy, breastfeeding</p> <p>Benzydamine - children under 12 years - oral rinse shouldn't be used.</p> <p>Lozenges- diabetes, children under 6 years (Strepsils®), children under 3 years (Throaties®, Tyrozets®), fructose intolerance, glucose-galactose malabsorption or sucrose-isomaltose insufficiency.</p> <p>Throat Sprays - children under 6 years, epiglottitis or methaemoglobinaemia.</p>	<p>Sore throat - NHS</p> <p>Sore Throat Patient.info</p>	<p>Sore throat - acute - NICE CKS</p>

Infrequent cold sores of the lip	Immunocompromised patients. 'Red flag' symptoms	Antiviral cold sore cream (aciclovir 5% cream)	Immunocompromised patients Terminally ill patients Only for treatment on face and lips	Cold sores - NHS Cold Sores Patient.info	Herpes simplex - oral - NICE CKS
Conjunctivitis	'Red flag' symptoms	Antimicrobial eye drops and eye ointment (chloramphenicol drops and ointment).	Children under 2 years Pregnancy Breastfeeding	Conjunctivitis - NHS Infective Conjunctivitis Patient.info	Conjunctivitis - infective - NICE CKS NICE Medicines evidence commentary: conjunctivitis and inappropriate prescribing. PHE Advice for schools: September 2017
Coughs and colds and nasal congestion	'Red flag' symptoms	Cough mixtures or linctus Saline nose drops Menthol vapour rubs Cold and flu capsules or sachets Analgesics Pseudoephedrine	Ibuprofen - long term conditions requiring regular pain relief; asthma; unstable high blood pressure; taking anticoagulants; stomach ulcer, perforation or bleeding (active or history of); renal, hepatic or cardiac impairment; pregnancy; breastfeeding; children under 3 months and not for use in chicken pox Paracetamol - long term conditions requiring regular pain relief, severe renal or severe hepatic impairment. Children under 2 months, babies born before 37 weeks and babies weighing less than 4 kg Pseudoephedrine - children under 12 years, pregnancy, breastfeeding, severe hypertension, severe coronary artery disease, taking MAOIs, severe renal impairment, diabetes, hyperthyroidism, closed angle glaucoma, phaeochromocytoma	Common cold - NHS PrescQIPP: Coughs and colds (Self care postcard - PrescQIPP login required) Common Cold Patient.info Coughs and Colds in Children Patient.info	Common cold - NICE CKS

Cradle cap (seborrhoeic dermatitis – infants)	If causing distress to the infant and not improving	Emulsifying ointment and Shampoos, including Capsal® Shampoo and Dentinox® Shampoo Cradle Cap	If it has spread to the face. Broken skin or infected skin	Cradle cap - NHS Cradle Cap Patient.info	Seborrhoeic dermatitis - NICE CKS
Haemorrhoids	'Red flag' symptoms	Haemorrhoid creams, ointments and suppositories.	Children under 18 years Pregnancy Breastfeeding Rectal bleeding or blood in the stool	Haemorrhoids (piles) - NHS Haemorrhoids Patient.info	Haemorrhoids - NICE CKS
Infant colic	'Red flag' symptoms	Simethicone and dimeticone suspensions Lactase drops	Dimeticone colic drops should be avoided in fructose, glucose-galactose or sucrose intolerant patients as it contains sucrose.	Colic - NHS PrescQIPP: Infant Colic (Self care postcard - PrescQIPP login required) Colic in Babies and Infants Patient.info	Colic - infantile - NICE CKS
Mild cystitis	'Red flag' symptoms	Sodium citrate or potassium citrate granules	Sodium citrate 4g sachets products – men, children under 16 years, diabetes, heart disease, hypertension, renal disease, pregnancy and breastfeeding Potassium citrate products - children under 6 years, renal disease/impairment, pregnancy and breastfeeding	Cystitis - NHS Cystitis in Women Patient.info	Urinary tract infection (lower) - women - NICE CKS

Minor conditions suitable for self-care					
Mild Irritant Dermatitis	No routine exceptions have been identified. See page 2 for general exceptions.	Emollients, Steroid creams, Crotamiton (e.g. Eurax®).	Children under 12 years with certain brands (however Eurax® is licensed for children over 3 years and Hc45® Hydrocortisone Cream is licensed for children over 10 years) Pregnancy Breastfeeding Broken or inflamed skin Use on the eyes, face or ano-genital region	Contact dermatitis - NHS Contact Dermatitis Patient.info	Dermatitis - contact - NICE CKS
Dandruff	No routine exceptions have been identified. See page 2 for general exceptions.	Antidandruff shampoos (e.g. coal tar or selenium) Antifungal shampoos (e.g. ketoconazole)	All products Pregnancy, breastfeeding Coal tar products Children under 12 years, infected open skin lesions and sore or acute pustular psoriasis Selenium sulphide children under 5 years Ketoconazole Children under 12 years	Dandruff - NHS Dandruff Patient.info	Seborrhoeic dermatitis - NICE CKS
Diarrhoea (adults)	No routine exceptions have been identified. See page 2 for general exceptions. THIS RECOMMENDATION DOES NOT APPLY TO CHILDREN	Loperamide 2mg capsules Rehydration sachets	Loperamide - inflammatory bowel disease, post bowel surgery, post pelvic radiation and colorectal cancer Oral rehydration sachets - liver or renal disease, patients on low potassium/sodium diets and patients with diabetes (Note Dioralyte® Relief can be given to diabetics)	Diarrhoea and vomiting - NHS Diarrhoea Patient.info	Diarrhoea - adult's assessment - NICE CKS
Dry eyes/sore tired eyes	No routine exceptions have been identified. See page 2 for general exceptions, e.g. pre-existing long term conditions affecting the eyes, more severe dry eye symptoms. (Self-care should only be	Eye drops for dry/sore, tired eyes e.g. hypromellose 0.3% eye drops and carbomers	Children under 18 years (carbomer) Pregnancy Breastfeeding (check individual brands)	Dry eyes - NHS Dry Eyes Patient.info	Dry eye syndrome - NICE CKS

	recommended in people with mild to moderate symptoms, without inflammation, unrelated to a long-term condition or essential medication, and not currently using drops for another eye health problem).				
Earwax	No routine exceptions have been identified. See page 2 for general exceptions (e.g. perforated ear drum)	Drops containing sodium bicarbonate, hydrogen peroxide, olive oil or almond oil.	None identified	Earwax build-up - NHS Earwax Patient.info	Earwax - NICE CKS
Excessive sweating (hyperhidrosis)	No routine exceptions have been identified. See page 2 for general exceptions.	Aluminium chloride sprays, roll-ons, solutions.	Broken or infected skin External use only	Excessive sweating (hyperhidrosis) - NHS Excessive Sweating Patient.info	Hyperhidrosis - NICE CKS
Head lice	No routine exceptions have been identified. See page 2 for general exceptions (e.g. current skin conditions of the scalp or side effects of a severe infestation)	Creams or lotions for head lice (e.g. malathion or dimeticone) Bug busting (conditioner and head lice combs)	Creams and lotions - children under 6 months, pregnancy and breastfeeding (some products), avoid use on broken or inflamed skin Conditioner and head lice combs – no restrictions	Head lice and nits - NHS PrescQIPP: Head lice (Self care postcard - PrescQIPP login required) Head Lice and Nits Patient.info Treatment and Prevention of head lice Patient.info	Head lice - NICE CKS

<p>Indigestion and heartburn</p>	<p>No routine exceptions have been identified. See page 2 for general exceptions.</p>	<p>Antacid tablets or liquids</p> <p>Ranitidine 75mg Tablets</p> <p>OTC proton pump inhibitors e.g. esomeprazole 20mg tablets.</p> <p>Sodium alginate, calcium carbonate or sodium bicarbonate liquids/tablets.</p>	<p>Ranitidine Children under 16 years, pregnancy, breastfeeding, peptic ulceration, people taking NSAIDs (these patients should not self-medicate with ranitidine 75 mg tablets but seek their doctor's advice before use)</p> <p>Proton pump inhibitors Children under 18 years or adults over 55 years with new or recently changed symptoms, pregnancy, breastfeeding, jaundice or liver disease</p> <p>Gaviscon® Children under 12 years, renal failure, hypercalcaemia, nephrocalcinosis, kidney stones</p> <p>Gaviscon® Infant Infants under 12 months, children over 2 years, babies born before 37 weeks, renal impairment</p>	<p>Indigestion - NHS</p> <p>PrescQIPP: Indigestion and Heartburn (Self care postcard - PrescQIPP login required)</p> <p>Indigestion (Dyspepsia) Patient.info</p> <p>Heartburn Patient.info</p>	<p>Dyspepsia - proven functional - NICE CKS</p>
<p>Infrequent constipation</p>	<p>No routine exceptions have been identified. See page 2 for general exceptions.</p> <p>Laxatives are not recommended for children unless they are prescribed by a GP (however self-care advise can be given on diet, fluid intake and exercise)</p>	<p>Bisacodyl tablets</p> <p>Senna</p> <p>Ispaghula Husk granules</p> <p>Lactulose solution</p>	<p>Bisacodyl Acute inflammatory bowel diseases, severe dehydration, pregnancy and breastfeeding, and children under 10 years.</p> <p>Senna Severe dehydration, children under 18 years, pregnancy and breastfeeding.</p> <p>Ispaghula husk granules Children under 6 years and diabetes.</p> <p>Lactulose Galactosaemia, gastro-intestinal obstruction, digestive perforation or risk of digestive perforation</p>	<p>Constipation - NHS</p> <p>Constipation Patient.info</p> <p>Constipation in Children Patient.info</p>	<p>Constipation - NICE CKS</p>

Infrequent migraine	No routine exceptions have been identified. See page 2 for general exceptions (e.g. those with severe or recurrent migraines should continue to seek advice from their GP)	Migraine tablets (e.g. Migralève®) Painkillers Anti-sickness tablets (prochlorperazine maleate – Buccastem M®)	For this indication: Ibuprofen Long term conditions requiring regular pain relief; asthma; unstable high blood pressure; taking anticoagulants; stomach ulcer, perforation or bleeding (active or history of); renal, hepatic or cardiac impairment; children under 12 years; pregnancy ; breastfeeding Paracetamol Long term conditions requiring regular pain relief, children under 12 years, severe renal or severe hepatic impairment Co-codamol (e.g. Migralève® Yellow tablets) Children under 12 years, severe liver disease, breastfeeding and pregnancy Prochlorperazine maleate Children under 18 years, impaired liver function, existing blood dyscrasias, epilepsy, Parkinson's Disease, prostatic hypertrophy, narrow angle glaucoma, pregnancy and breastfeeding	Migraine - NHS PrescQIPP: Infrequent Migraines (Self care postcard - PrescQIPP login required) Migraine Patient.info Migraine Treatment and Medication Patient.info	Migraine - NICE CKS
Insect bites and stings	No routine exceptions have been identified. See page 2 for general exceptions.	Steroid creams or creams for itching (e.g. mepyramine maleate 2% - Anthisan®)	Mepyramine maleate 2% Broken or infected skin and children under 2 years Steroid preparations See above under Mild Irritant Dermatitis	Insect bites and stings - NHS PrescQIPP: Insect bites and stings (Self care postcard - PrescQIPP login required)	Insect bites and stings - NICE CKS
Mild acne	No routine exceptions have been identified. See page 2 for general exceptions.	Benzoyl peroxide products	Pregnancy, breastfeeding Damaged skin Elderly Children under 12 years	Acne - NHS Acne Patient.info Acne Treatments Patient.info	Acne vulgaris - NICE CKS

Mild dry skin	See page 2 for general exceptions.	Emollient creams, ointments and lotions	Broken or infected skin Some preparations have age restrictions	Emollients - NHS	Eczema - atopic - NICE CKS
Sunburn due to excessive sun exposure	See page 2 for general exceptions (e.g. severe sunburn in babies and young children)	After sun products	No restrictions	Sunburn - NHS PrescQIPP: Sunburn (Self care postcard - PrescQIPP login required) Sunburn Patient.info	
Sun protection	ACBS approved indication of photodermatoses (i.e. where skin protection should be prescribed). See page 2 for general exceptions.	Sun protection cream, sprays, gels	No restrictions		
Mild to moderate hay fever/seasonal rhinitis	No routine exceptions have been identified. See page 2 for general exceptions.	Antihistamine tablets or liquids. Steroid nasal sprays Sodium cromoglicate eye drops	Chlorphenamine Tablets - children under 6 years 2mg/5ml solution - children under 1 year Taken monoamine oxidase inhibitors within the last 14 days; epilepsy; raised intra-ocular pressure including glaucoma; prostatic hypertrophy; severe hypertension or cardiovascular disease; bronchitis, bronchiectasis or asthma; hepatic impairment; renal impairment; rare hereditary problems of galactose intolerance, Lapp lactase deficiency or glucose-galactose malabsorption; pregnancy and breastfeeding. Loratadine Tablets - children under 6 years 1mg/ml syrup - children under 2 years	Hay fever - NHS PrescQIPP: Hay fever (Self care postcard - PrescQIPP login required) Hay Fever and Seasonal Allergies Patient.info	Allergic rhinitis - NICE CKS

			<p>Rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption; liver impairment; pregnancy and breastfeeding.</p> <p>Acrivastine Children under 12 years and adults over 65 years; renal impairment; rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption; pregnancy and breastfeeding.</p> <p>Cetirizine Tablets - children under 6 years 1mg/ml solution - children under 2 years (depending on the brand)</p> <p>Epilepsy; renal impairment; rare hereditary problems of fructose intolerance; pregnancy and breastfeeding.</p> <p>Beclomethasone 0.05% nasal spray-Children under 18 years, pregnancy and breastfeeding.</p> <p>Sodium Cromoglicate 2% eye drops Children under 6 years, pregnancy and breastfeeding.</p>		
Minor burns and scalds	<p>See page 2 for general exceptions. No routine exceptions have been identified.</p> <p>However more serious burns always require professional medical attention. Burns requiring hospital A&E treatment include but are not limited to: -all chemical and electrical burns; -large or deep burns;</p>	<p>Antiseptic burns cream or powder</p> <p>Cooling burn gel.</p>	<p>Betadine® Dry Powder Spray Pregnancy, breastfeeding, children under 2 years and taking lithium therapy</p>	<p>Burns and scalds - NHS</p> <p>Burns and Scalds Patient.info</p>	<p>Burns and scalds - NICE CKS</p>

	-burns that cause white or charred skin; -burns on the face, hands, arms, feet, legs or genitals that cause blisters.				
Minor conditions associated with pain, discomfort and/fever. (e.g. aches and sprains, headache, period pain, back pain)	No routine exceptions have been identified. See page 2 for general exceptions.	Paracetamol 500mg tablets Ibuprofen 200mg and 400mg tablets Co-codamol 8/500mg tablets NSAID topical creams or gels (e.g. Ibuprofen 5%gel) Paracetamol and Ibuprofen Suspension	Paracetamol Long term conditions requiring regular pain relief; severe renal or severe hepatic impairment; children under 2 months; babies born before 37 weeks; and babies weighing less than 4 kg. Ibuprofen Long term conditions requiring regular pain relief; asthma; unstable high blood pressure; severe renal, hepatic or cardiac impairment; taking anticoagulants; stomach ulcers/problems; children under 3 months; pregnancy; breastfeeding; and not for use in chicken pox Co-codamol Children under 12 years; severe liver disease; breastfeeding and pregnancy Ibuprofen 5% gel Children under 14 years; pregnancy and breastfeeding; broken/ infected skin	10 ways to reduce pain - NHS Back pain - NHS Fever in children - NHS Headaches - NHS Headaches Patient.info Period pain - NHS Period Pain (Dysmenorrhoea) Patient.info PrescQIPP: Pain (Self care postcard - PrescQIPP login required)	Analgesia - mild-to-moderate pain - NICE CKS
Mouth ulcers	No routine exceptions have been identified. See page 2 for general exceptions.	Lidocaine hydrochloride/ Chlorocresol/ Cetylpyridinium chloride e.g. Anbesol® Choline salicylate/Cetalkonium chloride e.g. Bonjela® Benzocaine e.g. Orajel®	Various age restrictions Active peptic ulceration (some preparations) Pregnancy Breastfeeding	Mouth ulcers - NHS Mouth Ulcers Patient.info	Aphthous ulcer - NICE CKS

Nappy rash	No routine exceptions have been identified. See page 2 for general exceptions.	Nappy rash creams (e.g. Metanium®, Sudocrem®)	No restrictions	Nappy rash - NHS Nappy Rash Patient.info	Nappy rash - NICE CKS
Oral thrush	No routine exceptions have been identified. See page 2 for general exceptions.	Miconazole 2% oral gel (Daktarin® oral gel)	Children under 4 months Pregnancy, breastfeeding Patients taking warfarin Liver dysfunction	Oral thrush (mouth thrush) - NHS Oral thrush in babies - NHS	Candida - oral - NICE CKS
Prevention of dental caries	No routine exceptions have been identified. See page 2 for general exceptions. Note that some fluoride toothpastes (2800ppm and 5000ppm) are POM Prescription only medicines should be re-directed to the dentist to prescribe	Fluoride toothpastes Mouthwashes such as chlorhexidine gluconate	Children under 12 years	Tooth decay - NHS Tooth Decay Patient.info	
Ringworm/ Athlete's foot	Lymphoedema or history of lower limb cellulitis. See page 2 for general exceptions.	Antifungal creams or sprays (e.g. miconazole or terbinafine)	All products Diabetes, pregnancy, breastfeeding. Terbinafine hydrochloride Children under 16 years Some preparations may only be licensed for athlete's foot.	Athlete's foot - NHS Ringworm - NHS	Fungal skin infection - foot - NICE CKS
Teething/mild toothache	No routine exceptions have been identified. See page 2 for general exceptions.	Teething gel (Calgel® and Dentinox®) Painkillers	Calgel® and Dentinox® teething gel Babies under 5 months, heart disease and liver disease Ibuprofen Long term conditions requiring regular pain relief; asthma; unstable high blood pressure; taking anticoagulants; stomach ulcer, perforation or bleeding (active or history of); renal, hepatic or cardiac impairment; pregnancy; breastfeeding; children under 3 months; and not for use in chicken pox.	Toothache - NHS Teething Patient.info PrescQIPP: Teething (Self care postcard - PrescQIPP login)	Teething - NICE CKS

			<p>Paracetamol Long term conditions requiring regular pain relief; severe renal or severe hepatic impairment; children under 2 months; babies born before 37 weeks; and babies weighing less than 4 kg.</p>	<p>required) PrescQIPP: Toothache (Self care postcard - PrescQIPP login required)</p>	
Threadworms	No routine exceptions have been identified. See page 2 for general exceptions.	Mebendazole 100mg tablets	Children under 2 years Pregnancy Breastfeeding	Threadworms - NHS	Threadworm - NICE CKS
Travel Sickness	No routine exceptions have been identified. See page 2 for general exceptions	Travel sickness tablets	<p>Cinnarizine- Parkinson's Disease; fructose or galactose intolerance, Lapp lactase deficiency, glucose-galactose malabsorption or sucrase-isomaltase insufficiency; porphyria; children under 5 years; pregnancy and breastfeeding.</p> <p>Hyoscine- Children under 3 years (Joyrides®); under 4 years (Kwells®); epilepsy; glaucoma; pregnancy and breastfeeding; paralytic ileus; pyloric stenosis; prostatic enlargement; myasthenia gravis.</p>	<p>Motion sickness - NHS</p> <p>Patient info: Motion Sickness</p>	
Warts and Verrucae	No routine exceptions have been identified. See page 2 for general exceptions	<p>Gels, skin paints, ointments and medicated plasters containing salicylic acid and lactic acid</p> <p>Wart freeze treatments (e.g. Dimethyl ether propane cold spray)</p>	<p>Diabetic patients Face, neck, ano-genital area Moles, birthmarks, hairy warts or skin lesions Broken skin Impaired peripheral blood circulation</p> <p>Age restrictions apply to some preparations (e.g. wart freeze treatments should not to be used in children under 4 years)</p>	<p>Warts and verrucas - NHS</p> <p>Warts and Verrucas Patient.info</p>	Warts and verrucae - NICE CKS

*This list may not be exhaustive. If needed see summary of product characteristics and patient information leaflets within the [EMC](#)

Useful resources

NHS Barnsley CCG 'Prescribing of over the counter medicines is changing' posters and leaflets to support implementation can be obtained from the Medicines Management Team. The following may also be of use:

NHS England Implementation Resources:

<https://www.england.nhs.uk/medicines/conditions-for-which-over-the-counter-items-should-not-routinely-be-prescribed/conditions-for-which-over-the-counter-items-should-not-routinely-be-prescribed-in-primary-care-implementation-resources/>

Resources for CCGs and healthcare professionals on the implementation of not routinely prescribing over the counter (OTC) medicines for certain conditions.

Quick reference guide for healthcare professionals: <https://www.england.nhs.uk/publication/quick-reference-guide-for-healthcare-professionals-conditions-for-which-over-the-counter-items-should-not-routinely-be-prescribed-in-primary-care/>

Patient information leaflet: <https://www.england.nhs.uk/wp-content/uploads/2018/08/1a-over-the-counter-leaflet-v1.pdf>

Patient information leaflet for print: <https://www.england.nhs.uk/wp-content/uploads/2018/08/1a-over-the-counter-leaflet-print-ready.pdf>

Easy to read patient information leaflet: <https://www.england.nhs.uk/wp-content/uploads/2018/08/1b-easy-read-over-the-counter-leaflet.pdf>

Patient Information Sheet (space to add notes/specific advice as needed): <https://www.england.nhs.uk/wp-content/uploads/2018/08/1c-over-the-counter-patient-info-sheet.pdf>

PrescQIPP Self-Care Resources (PrescQIPP login required):

<https://www.prescqipp.info/our-resources/webkits/self-care/>

These resources focus on reducing the prescribing of over the counter preparations in general practice, signposting to the correct health service and supporting the implementation of the NHS England OTC guidance. Each set of resources contains tools that can be adapted for local use before implementation.

Coughs and colds poster: <https://www.prescqipp.info/media/3862/cough-colds-and-sore-throats-poster.jpg>

Medicines cabinet poster: <https://www.prescqipp.info/media/3756/medicine-cabinet-poster-v1-l.jpg>

Choose self-care poster: <https://www.prescqipp.info/media/3754/feeling-under-the-weather-poster-v1-l.jpg>

'No prescription required' leaflet: <https://www.prescqipp.info/our-resources/webkits/self-care/resources-for-professionals/>

Receptionist information sheet checklist: <https://www.prescqipp.info/our-resources/webkits/self-care/resources-for-professionals/>

Self-Care Forum:

<http://www.selfcareforum.org/>

Aims to raise the awareness of self-care and embed it into everyday life. The website contains lots of useful resources and information for prescribers and patients.

NHS.uk:

<https://www.nhs.uk/>

Aims to help patients take control of their health and wellbeing

The Royal Pharmaceutical Society:

<https://www.nhs.uk/live-well/healthy-body/your-medicine-cabinet/>

The RPS offers advice on over the counter products that should be kept in a medicine cabinet at home to help patients treat a range of self-treatable illnesses.

References

1. NHS England: Conditions for which over the counter items should not routinely be prescribed in primary care: Guidance for CCGs. Available at: <https://www.england.nhs.uk/wp-content/uploads/2018/03/otc-guidance-for-ccgs.pdf> Accessed 22.01.19
2. PrescQIPP CIC Bulletin 227: Over the counter items – GP guide to self-care. Available at: <https://www.prescqipp.info/our-resources/bulletins/bulletin-227-self-care-over-the-counter-items/favicon.ico> Accessed 22.01.19
3. PrescQIPP self-care resources. Available at: <https://www.prescqipp.info/our-resources/webkits/self-care/> Accessed 22.01.19
4. Electronic Medicines Compendium. Available at: <https://www.medicines.org.uk/emc/> Accessed 22.01.19
5. NHS.uk Website. Available at: <https://www.nhs.uk/conditions/> Accessed 22.01.19
6. Patient.info Website. Available at: <https://patient.info/> Accessed 26.3.19
7. CQC Treating minor ailments and promoting self-care in adult social care. Available at: <https://www.cqc.org.uk/guidance-providers/adult-social-care/treating-minor-ailments-promoting-self-care-adult-social-care> Accessed 26.3.19

Acknowledgements

Sheffield CCG Self-Care Guidance, October 2018. Available at:

https://www.intranet.sheffieldccg.nhs.uk/Downloads/Medicines%20Management/prescribing%20guidelines/Self-care/Self-care_guidance.pdf

Leeds CCG Implementing the NHSE guidance in care settings. Available at: <https://www.leedscg.nhs.uk/health/campaigns/over-the-counter-medicines-changes-to-prescribing/>

Development Process

This guidance was approved by the Barnsley Area Prescribing Committee on 12th June 2019